

AD-A071 060

ROCHESTER UNIV NY DEPT OF STATISTICS  
THE LOGIT METHOD FOR COMBINING TESTS.(U)  
1979 E O GEORGE, G S MUDHOLKAR

F/G 12/1

UNCLASSIFIED

AFOSR-TR-79-0742

AFOSR-77-3360


NL

1 OF 1  
AD  
A071060



END  
DATE  
FILMED

8-79  
DDC



NATIONAL BUREAU OF STANDARDS  
MICROCOPY RESOLUTION TEST CHART

THE LOGIT METHOD FOR  
COMBINING TESTS

1979

by

E. OLUSEGUN GEORGE and GOVIND S. MUDHOLKAR \*

## A B S T R A C T

A new combination statistic, the sum of the logits of P-values is introduced and its standardized form is found to be essentially equivalent to a standardized Student t in null distribution. Comparisons with Fisher's omnibus procedure by Bahadur ARE, membership in a complete class of tests and a Monte Carlo simulation of the power functions, show that both procedures are equivalent by the first criterion and neither procedure is generally dominant by ~~either~~ either of the two criteria.


Key words: Combining tests, Logit, Fisher's Omnibus procedure, Bahadur Efficiency, admissibility.

\*E. Olusegun George is lecturer in mathematics and statistics at the University of Ife, Ile-Ife, Nigeria. Govind S. Mudholkar is professor of statistics and biostatistics, University of Rochester, Rochester NY 14627. This research was sponsored in part by the Air Force Office of Scientific Research, Air Force Systems Command, USAF under Grant No. AFOSR-77-3360 and by the African American Institute.

Approved for public release;  
distribution unlimited.

AD A071060

DDC FILE COPY



## 1. INTRODUCTION

Several authors have studied the problem of combining independent tests, especially noteworthy among whom are Fisher (1932), Karl Pearson (1933) E.S. Pearson (1938), Wallis (1942), A. Birnbaum (1954), Liptak (1958), van Zwet and Oosterhoff (1967), Oosterhoff (1969), and Littell and Folks (1971, 1973). The combination problem has been cast and analyzed in various canonical forms by these authors, but for the purposes of this essay its essentials may be described as follows: Let  $T_i$ ,  $i=1,2,\dots,k$  be  $k$  independently distributed statistics, the large values of which are significant in testing respective null hypotheses  $H_{i0}: \theta_i = \theta_{i0}$  against one-sided alternatives  $H_{i1}: \theta_i > \theta_{i0}$  concerning real-valued parameters  $\theta_i$  of the distributions of  $T_i$ ,  $i=1,2,\dots,k$ . The problem is to find a reasonable combination, i.e. a function of  $T_1, T_2, \dots, T_k$  which may be used for testing the overall hypothesis  $H_0: \theta_i = \theta_{i0}$ ,  $i=1,2,\dots,k$  versus the alternative  $H_1: \theta_i \geq \theta_{i0}$ ,  $i=1,2,\dots,k$  with at least one inequality strict. Many problems of combining tests of composite hypotheses, such as F-test of the general linear hypothesis, can be reduced to this form.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)  
NOTICE OF TRANSMITTAL TO RDC  
This technical report has been reviewed and is  
approved for public release in accordance with AFR 190-12 (7b).  
Distribution is unlimited.  
A. D. BLOSE  
Technical Information Officer



In describing the combination problem it is often convenient to use the P-values  $P_i$  of the tests as the pivotal entities instead of the statistics  $T_i$ ; a combination statistic is then a function of  $P_1, P_2, \dots, P_k$ . The P-values  $P_i$  is of course the probability, under the null hypothesis, of obtaining at least as extreme a value of  $T_i$  as observed. Thus denoting the null distribution of  $T_i$  by  $F_{i0}$  the P-value is given by

$$\begin{aligned} P_i &= 1 - F_{i0}(T_i), & \text{if large values of } T_i \text{ are significant} \\ &= F_{i0}(T_i), & \text{if small values of } T_i \text{ are significant.} \end{aligned} \quad (1.1)$$

An advantage of using  $P_i$  is that it can be interpreted by itself, without a reference to the distribution of  $T_i$ . Moreover, if  $F_{i0}$  is continuous the null distribution of  $P_i$  is uniform on  $(0,1)$ . This latter property of the P-values makes the null distributions of combination statistic manageable. In fact, most common combination statistics in the literature have simple null distributions. Notable examples of such statistics are (i)  $T^{(T)} = \min P_i$ , due to Tippett (1931); (ii)  $T^{(W)} = P_{(r)}$ , the  $r^{\text{th}}$  largest of the  $k$  P-values, due to Wilkinson (1951); (iii)  $T^{(F)} = -2 \sum \log P_i$ , due to Fisher; (iv)  $T^{(P)} = -2 \sum \log(1 - P_i)$ , due to Pearson; and (v)  $T^{(N)} = \sum \alpha_i \Phi^{-1}(1 - P_i)$ , due to Liptak, where  $\Phi(\cdot)$  is the standard normal d.f. and

$\alpha_i \geq 0$  are arbitrary weights. Under the overall null hypothesis  $H_0$ ,  $T^{(T)}$  and  $T^{(W)}$  have beta distributions,  $T^{(F)}$  and  $T^{(P)}$  are  $\chi^2_{2k}$ -variates and the Liptak's statistic  $T^{(L)}$  is normally distributed. Among these procedures Fisher's procedure, which rejects  $H_0$  when  $T^{(F)}$  is large, has been shown by Littell and Folks (1973) to be most efficient with respect to Bahadur's measure of A.R.E. A fortiori, Littell and Folks (1971, 1973) demonstrated that Fisher's method has maximum Bahadur A.R.E. among all reasonable combination methods based upon the P-values.

Furthermore Birnbaum (1954) proved that Wilkinson's (1951) and Pearson's (1933) method are inadmissible and recommended the use of Fisher's method for combining statistics whose distributions belong to the exponential family.

In this paper we introduce a new combination statistic, the logit statistic  $T_L = -\sum_{i=1}^k \log [P_i / (1-P_i)]$  and suggest rejecting  $H_0$  when it is large. We show that this procedure has the same Bahadur efficiency as Fisher's method and consequently, because of a result of Littell and Folks (1973), both procedures are optimal.

Accession For	NTIS GRA&I	<input checked="" type="checkbox"/>
	DOC TAB	<input type="checkbox"/>
	Unannounced	<input type="checkbox"/>
	Justification	<input type="checkbox"/>
Distribution/		
Availability Codes		
Avail and/or	Special	
Dist		

The Logit statistic, its exact null distribution and a simple t-approximation to the null distribution are given in section 2. In section 3 the exact Bahadur slope is computed. In section 4 we answer a question raised by Oosterhoff (1969, p.42) on the admissibility of Fisher's method and discuss some complete class results for both methods. In section 5 we describe a Monte Carlo simulation of the power functions of the procedures when combining Student t tests.

## 2. THE LOGIT PROCEDURE

Following Berkson's (1944) term for the log-odds ratio  $\log[P/(1-P)]$  the combination statistic

$$T_L = -\sum_{i=1}^k \log[P_i/(1-P_i)] \quad (2.1)$$

is termed the Logit statistic. When the null distribution  $F_{10}$  of  $T_1$  is continuous  $P_1$  is uniformly distributed on  $(0,1)$ . Consequently under the null hypothesis  $H_{10} - \log[P_1/(1-P_1)]$  is distributed according to the logistic distribution function

$$F(z) = [1 + \exp(-z)]^{-1}, \quad -\infty < z < \infty \quad (2.2)$$



and the Logit statistic is a sum of i.i.d. logistic variates under  $H_0$ . George and Mudholkar (1977) have shown that the exact null distribution,  $F_0$  of  $T_L$ , is given by

$$1-F_0(z) = \sum_{p=0}^{k-1} \sum_{r=0}^{k-1-p} \sum_{m=1}^{p+r+1} (-1)^{r+1} A_{k,p} S_{p+r+1,m} \frac{z^r}{r} \binom{k-1}{p+r} (m-1)! \\ [e^{-z}/(1-e^{-z})]^m \quad (2.3)$$

when  $k$  is even, and by

$$1-F_0(z) = \sum_{p=0}^{k-1} \sum_{r=0}^{k-1-p} \sum_{m=1}^{p+r+1} (-1)^{r+m+1} A_{k,p} S_{p+r+1,m} \frac{z^r}{r} \binom{k-1}{p+r} (m-1)! \\ [e^{-z} (1+e^{-z})]^k \quad (2.4)$$

when  $k$  is odd; where  $A_{k,p}$ 's are computed from the following equations.

$$[\pi s / \sin \pi s]^k = A_{k,0} + A_{k,1}s + A_{k,2}s^2 + \dots \quad (2.5)$$

and

$$(\pi s / \sin \pi s) = \sum_{m=0}^{\infty} (-1)^{m-1} [(2^{2m}-2)/(2m)!] B_{2m}(\pi s) 2^m, \quad (2.6)$$

where  $B_{2m}$ 's are Bernoulli numbers, and  $S_{n,m}$ 's are Stirling numbers of the second kind. Thus when  $k = 2$ ,

$$1-F_2(z) = z[e^{-z}/(1-e^{-z})]^2 + (z-1)[e^{-z}/(1-e^{-z})] \quad (2.7)$$

and when  $k = 3$

$$1-F_3(z) = e^{-z}/(1+e^{-z}) - 2ze^{-z}/(1+e^{-z})^2 \\ + (z^2 + \pi^2)e^{-z}(1+e^{-z})/2(1+e^{-z})^3 . \quad (2.8)$$

George and Mudholkar (1977) have illustrated that the convolution of  $k$ -logistic random variables, and hence the null distribution of the Logit statistic, is well approximated by the normal distribution with equal variance, that this approximation becomes very good when Edgeworth-corrected, and that a multiple of the student's  $t$ -distribution with the degrees of freedom obtained by considering the kurtosis provides a simple and an even better approximation. Specifically, for the null distribution of Logit statistic, it was proposed that

$$-\sum_{i=1}^k \log[P_i/(1-P_i)] \approx \frac{k(5k+2)}{3(5k+4)} t_{5k+4} \quad (2.9)$$

In Table 1 the quality of this approximation is illustrated numerically for  $k=3$ .



### 3. BAHADUR A.R.E. OF THE LOGIT STATISTIC

The concept of Bahadur A.R.E. is well discussed in the literature (Bahadur 1971). Let  $T_n$  be a statistic for testing a null hypothesis  $H_0: \theta \in \mathcal{H}_0$  against an alternative  $H: \theta \in \mathcal{H}_1$ . Assume, without loss of generality, that large values of  $T_n$  are significant, and let  $T_n = t_n$  be observed. Then the rate of convergence to zero of the P-value  $L_n(t_n) = \Pr\{T_n \geq t_n | H_0\}$  as  $n \rightarrow \infty$  is a measure of the efficiency of the test  $T_n$ . Specifically suppose that for  $i = 1, 2$  and  $\theta \in \mathcal{H}_1$ , there exist positive numbers  $c_i(\theta)$  such that for sequences of tests  $\{T_n^{(i)}\}$  with corresponding P-values  $\{L_n^{(i)}\}$

$$\lim_{n \rightarrow \infty} -\frac{1}{n} \log L_n^{(i)} = c_i(\theta) \text{ a.s. } [P_\theta] \quad (3.1)$$

Then the Bahadur A.R.E. of  $\{T_n^{(1)}\}$  relative to  $\{T_n^{(2)}\}$  is given by

$$\Psi_{12}(\theta) = c_1(\theta)/c_2(\theta) \quad (3.2)$$

Since a Bahadur slope is not always easy to compute, Bahadur (1971) gave the following result to facilitate its computation.

Theorem (3.1) (Bahadur). Suppose that

$$n^{-1/2} T_n \rightarrow b(\theta) \text{ a.s. } P_\theta, \theta \in \mathcal{H}_1, \quad (3.3)$$

where  $-\infty < b(\theta) < \infty$  and

$$\lim_{n \rightarrow \infty} n^{-1} \log[1 - F_n(\sqrt{n}t)] = -f(t) \quad (3.4)$$

for each  $t$  in some open interval  $I$ , where  $F_n(\cdot)$  is the null distribution function of  $T_n$ ,  $f$  is continuous on  $I$  and  $\{b(\theta), \theta \in \mathcal{H}_1\} \subset I$ . Then

$$\lim_{n \rightarrow \infty} n^{-1} \log L_n \text{ exists}$$

and

$$c(\theta) = 2f(b(\theta)). \quad (3.5)$$

Theorem 3.2: For  $i = 1, \dots, k$ , let  $\{T_{n_i}^{(1)}\}$  be sequences of statistics for respectively testing  $H_{i0}: \theta_i = \theta_{i0}$  against  $H_{i1}: \theta_i > \theta_{i0}$ , and let  $\{L_{n_i}^{(1)}\}$  be the corresponding  $F$ -values. Assume that the  $T_{n_i}^{(1)}$ 's are independently distributed and that there exists a positive value function  $c_i(\theta_i)$  defined for  $\theta_i > \theta_{i0}$  such that

$$\lim_{n_i \rightarrow \infty} n_i^{-1} \log L_{n_i}^{(i)} = c_i(\theta_i)/2 \quad \text{a.s.} \quad (3.6)$$

Also assume that

$$\lim_{n_i \rightarrow \infty} n_i/n = \lambda_i, \quad (3.7)$$

where  $n_k = n_1 + \dots + n_k$ . Then the exact slope of the Logit combination procedure for testing  $H_0: \theta_i = \theta_{i0}$ ,  $i = 1, \dots, k$  against  $H_1: \theta_i \geq \theta_{i0}$ ,  $i=1, \dots, k$  with at least one strict inequality, is given by

$$c_L(\theta_1, \dots, \theta_k) = \sum_{i=1}^k \lambda_i c_i(\theta_i). \quad (3.8)$$

Proof: It is well known that if  $\phi_n(T_n)$  is a strictly increasing and continuous function of  $T_n$ , then  $\{\phi_n(T_n)\}$  and  $\{T_n\}$  have the same exact slopes.

Hence let

$$\begin{aligned} T_n^{(L)*} &= n^{-1/2} T_n^{(L)} \\ &= n^{-1/2} \sum_{i=1}^k \log[(1 - L_{n_i}^{(i)})/L_{n_i}^{(i)}] \end{aligned}$$

Then

$$\begin{aligned} n^{-1/2} T_n^{(L)\star} &= \sum_{i=1}^k n^{-1} \log(1 - L_{n_i}^{(i)}) - \sum_{i=1}^k n^{-1} \log L_{n_i}^{(i)} \\ &= \sum_{i=1}^k (n_i/n) n_i^{-1} \log(1 - L_{n_i}^{(i)}) - \sum_{i=1}^k (n_i/n) n_i^{-1} \log L_{n_i}^{(i)}. \end{aligned}$$

Clearly  $\lim_{n_i \rightarrow \infty} n_i^{-1} \log L_{n_i}^{(i)} = c_i(\theta_i)$  w.p.1 implies that

$$\lim_{n_i \rightarrow \infty} n_i^{-1} \log(1 - L_{n_i}^{(i)}) = 0 \text{ w.p.1.}$$

Hence from (3.6) and (3.7), it follows that

$$\lim_{n \rightarrow \infty} n^{-1} 2 T_n^{(L)\star} = \sum_{i=1}^k \lambda_i c_i(\theta_i) \text{ w.p.1}$$

Now let  $F_0^\star$  denote the d.f of  $T_n^{(L)\star}$ . Then

$$1 - F_0^\star(t) = 1 - F_0(\sqrt{n}t) \quad -\infty < t < \infty, \quad (3.9)$$

where  $F_0$  is the d.f. of  $T_n^{(L)}$ . By combining the expressions for d.f. of  $F_0$  given by (2.3) and (2.4), it can easily be shown that

$$\begin{aligned} 1 - F_0^\star(\sqrt{n}z) &= [e^{-nz}/(1 - e^{-nz})] h_1(nz) \quad \text{if } k \text{ is even,} \\ &= [e^{-nz}/(1 + e^{-nz})] h_2(nz) \quad \text{if } k \text{ is odd,} \end{aligned} \quad (3.10)$$



where for  $j = 1, 2$

$$h_j(z) = 1 + p_{1,j}(z) [e^{-z} / (1 + (-1)^j e^{-z})]^{r_{1,j}} + \dots \\ + p_{s,j}(z) [e^{-z} / (1 + (-1)^j e^{-z})]^{r_{s,j}} ,$$

$p_{r,j}(z)$ ,  $1 \leq r \leq s$ , are polynomials in  $z$ ,

and  $r_{m,j,s}$ ,  $1 \leq r_{m,j} \leq k$ , are positive integers.

From (3.10) we get

$$-\frac{1}{n} \log[1 - F_0^*(\sqrt{n} z)] = -z - \frac{1}{n} (\log(1 - e^{-nz}) + \log h_1(nz)) \text{ if } k \text{ is even,} \\ = -z - \frac{1}{n} (\log(1 + e^{-nz}) + \log h_2(nz)) \text{ if } k \text{ is odd.}$$

Clearly  $\lim_{n \rightarrow \infty} h_j(nz) = 1$ . Hence

$$-\lim_{n \rightarrow \infty} \frac{1}{n} \log[1 - F_0^*(\sqrt{n} z)] = z.$$

Using Theorem 3.1, we immediately get the exact slope of

$T_n^{(L)*}$  to be  $\sum_{i=1}^k \lambda_i c_i(\theta_i)$ . Littell and Folks (1971) have shown

that under the conditions of Theorem 3.2, the exact slope of Fisher's procedure is  $\sum_{i=1}^k \lambda_i c_i(\theta_i)$ . Furthermore they showed (1973) that among "essentially" all combination procedures



based on P-values, Fisher's procedure is optimal according to Bahadur A.R.E. We state a slightly more general version of this result, which holds for the Logit and Fisher's procedures.

Theorem 3.3: Let  $T_n$  be a combination statistic for testing  $H_0$  against  $H_1$  based on P-values  $L_{n_1}^{(1)}, \dots, L_{n_k}^{(k)}$ . Suppose that  $T_n$  satisfies the following conditions.

I.  $T_n$  is non-decreasing in each  $L_{n_i}^{(i)}$ , i.e.,

$$l'_1 \leq l_1, \dots, l'_k \leq l_k \text{ imply } T_n(l'_1, \dots, l'_k) \leq T_n(l_1, \dots, l_k).$$

In this case small values of  $T_n$  are significant.

II.  $T_n$  is non-increasing in each  $L_{n_i}^{(i)}$ , i.e.,

$$l_1 \leq l'_1, \dots, l_k \leq l'_k \text{ imply } T_n(l_1, \dots, l_k) \geq T_n(l'_1, \dots, l'_k).$$

In this case large values of  $T_n$  are significant.

Then the slopes  $c(\theta_1, \dots, \theta_k)$  of  $\{T_n\}$  and  $c_L(\theta_1, \dots, \theta_k)$  of the Logit and Fisher's procedures satisfy the inequality  $c(\theta_1, \dots, \theta_k) \leq c_L(\theta_1, \dots, \theta_k)$ .

REMARK: Wieand (1976) has discussed conditions under which Bahadur's and Pitman's methods of comparing tests coincide. Under the conditions he stated both the Logit and Fisher's methods of combining tests are optimal with respect to Pitman's efficiency as well.

TABLE 1

THE EXACT AND APPROXIMATE \*D.F.  
OF STANDARDIZED LOGIT STATISTIC,  $k=3$

x	Exact	Approximate	Error
.05	.5209	.5208	0.0001
.25	.6033	.6028	0.0005
.45	.6809	.6802	0.0007
.65	.7506	.7499	0.0007
1.00	.8486	.8482	0.0004
1.20	.8903	.8901	0.0002
1.45	.9291	.9291	0.0000
1.75	.9598	.9600	-0.0002
2.50	.9918	.9920	-0.0002
3.00	.9975	.9975	0.0000

\*The approximation is based upon Equation 2.9.

#### 4. ADMISSIBILITY AND COMPLETE CLASS RESULTS

Several researchers (A. Birnbaum 1954, 1955; van Zwet and Oosterhoff 1967; Oosterhoff 1969 and Brown, Cohen and Strawderman 1976) have discussed the question of admissibility and membership in complete classes of combination procedures. Birnbaum (1955) proved that if the distributions of the component test statistics  $T_i$  are one-parameter exponential families, a procedure for combining these statistics can be admissible only if its acceptance region is monotone and convex in the  $(t_1, \dots, t_k)$  space. He further showed that this condition is both necessary and sufficient for admissibility if for each  $i$ ,  $T_i$  is normally distributed with a known variance. Oosterhoff considered the more general problem in which the distribution of the  $T_i$ 's have strict monotone likelihood ratio (MLR) properties, then monotone procedures are essentially complete. Brown et al showed that under the conditions stated by Oosterhoff monotone procedures are complete. We now compare the Logit and Fisher's procedures in the light of the above results.

First we answer a question raised by Oosterhoff (1969, page 42) and show that Fisher's procedure is admissible

when the distributions of the component statistics are one-parameter exponential families.

Theorem 4.1: If for each  $i$ , the statistic  $T_i$  for testing  $H_{i0}: \theta_i = \theta_{i0}$  against  $H_{i1}: \theta_i > \theta_{i0}$  have densities that can be expressed in the canonical form

$$f_i(t_i) = h(t_i) c(\theta_i) e^{\theta_i t_i} \quad (4.1)$$

$i = 1, \dots, k$  and if the  $T_i$ 's are independent, then Fisher's combination procedure is admissible for testing  $H_0: \theta_i = \theta_{i0}, i=1, \dots, k$  against  $H_1: \theta_i \geq \theta_{i0}, i=1, \dots, k$  with at least one inequality strict.

The following result given by Mudholkar (1969) is used in the proof.

Lemma 4.1: Let  $g_i(x_i)$  be positive functions of  $x_i$ ,  $i=1, \dots, k$ . If  $g_i(x_i)$  is a logconcave function of  $x_i$  for each  $i$ , then the set  $A = \{(x_1, \dots, x_k): \prod_{i=1}^k g_i(x_i) > c\}$  is convex for any positive real constant  $c$ .

Proof of theorem 4.1: It is easily shown that if the distribution of  $T_i$  is given by equation 4.1, then P-value corresponding to observed value  $T_i = t_i$  is given by

$$P_i(t_i) = 1 - F_{i0}(t_i).$$



Consequently, the acceptance region

$$A_F = \{(t_1, \dots, t_k) : -2 \sum_{i=1}^k \log P_i(t_i) < c_F\},$$

where  $c_F$  is a constant, of the Fisher's procedure, may be expressed as

$$A_F = \{(t_1, \dots, t_k) : \prod_{i=1}^k (1 - F_{i0}(t_i)) > e^{-c_F/2}\}$$

Now for any vector  $(t'_1, \dots, t'_k)$  such that  $t'_i < t_i$ ,  $i=1, \dots, k$ , it is clear that  $(t_1, \dots, t_k) \in A_F$  implies that

$(t'_1, \dots, t'_k) \in A_F$ . Hence  $A_F$  is a monotone acceptance region.

Now since  $T_i$  has an exponential family distribution,

$1 - F_{i0}(t_i)$  is logconcave. Therefore by Lemma 4.1,

$A_F$  is convex. The result of Birnbaum (1955) then implies that Fisher's procedure is admissible in the above context.

It is clear that in the exponential family case the acceptance region

$$A_L = \{(t_1, \dots, t_k) : \sum_{i=1}^k \log[(1 - F_{i0}(t_i))/F_{i0}(t_i)] > c_L\}$$

where  $c_L$  is a constant is not convex, however, convexity of an acceptance region is not known to be sufficient for admissibility.

Consequently the Logit procedure cannot be ruled out as inadmissible in the context of exponential family of distributions. Furthermore, in the larger context of



combining tests whose component statistics have M.L.R. or strict M.L.R. properties, results due to Brown et al (1976) can be used to show that both procedures belong to essentially complete or complete classes of tests. The combination of independent Student t test and independent F tests are particular cases of the M.L.R. families that are of practical importance.

We report here an empirical evidence that shows that in combining independent Student t tests, neither of the Logit or Fisher's procedures dominates the other uniformly.

## 5. A STUDY OF THE POWER FUNCTIONS BY SIMULATION

In this section the power functions of the Logit and Fisher's combination statistics are studied in the context combining Student t tests. Let  $t_{n_i-1}$ ,  $i=1,2$  be two independent Student t statistics with degrees of freedom  $n_1-1$  and  $n_2-1$  respectively. Let  $t_{n_1-1}$  be used to test  $H_{10}: \mu_1=0$  versus  $H_{11}: \mu_1 > 0$ , where  $\mu_1$  is the unknown mean of a normal population with unknown variance. Consider the problem of testing the combined null hypothesis  $H_0: \mu_1 = \mu_2 = 0$  against the alternative  $H_1$ : Either  $\mu_1 > 0$  or  $\mu_2 > 0$  by using the Logit statistic  $T^{(L)} = \log [(1-P_1)/P_1] + \log [(1-P_2)/P_2]$  or the Fisher's statistic  $T^{(F)} = -2 \log P_1 - 2 \log P_2$ , where  $P_1$  and  $P_2$  are the P-values corresponding to  $t_{n_1-1}$  and  $t_{n_2-1}$  respectively.

Although both  $T^{(L)}$  and  $T^{(F)}$  have simple null distributions under  $H_0$ , their distributions under  $H_1$  are complex, consequently it is futile to attempt to obtain their exact power functions. A reasonable estimates of the powers may be obtained from a Monte-Carlo experiment.

### 5.1 THE MONTE CARLO EXPERIMENT

For  $i=1,2$   $n_i$  independent standard normal variates are generated on the IBM 360/365 computer at the University of Rochester by using the McGill University random number package developed by a technique due to Marsaglia (1961) for generating standard normal deviates.  $N(\mu_i, 1)$  deviates are obtained by adding  $\mu_i$  to each generated variable. From these independent Student t statistics and their corresponding P-values are obtained.

The power functions of the Logit and Fisher's procedures are approximated from these statistics at levels of significance  $\alpha=.01$  and  $.05$  and for sample sizes  $n_1=n_2=5$  and  $n_1=n_2=10$ . The power of each procedure is estimated at  $\mu_1=0.0$  (0.1) 1.6 and  $\mu_2 = 0.0$  (0.1) 1.6 by the proportion of times the procedure rejects  $H_0$  in 3000 trials at each  $(\mu_1, \mu_2)$ .

### 5.2 RESULTS

Table 2 represents some estimates of the power functions of the Logit and Fisher's procedures. From the table it can be observed that the power of each procedure is ncreasing in  $\mu_2$  when  $\mu_1$  is held fixed and vice-versa; and that the Logit procedure is slightly superior to Fisher's procedure around the equiangular line  $\mu_1=\mu_2$  while Fisher's procedure is superior elsewhere.

POWER FUNCTION OF THE LOGIT AND FISHER'S  
METHODS FROM A MONTE CARLO EXPERIMENT

$\mu_2$	$\mu_1$								
	0.0	0.2	0.4	0.6	0.8	1.0	1.2	1.4	1.6
$\alpha = .05, n_1 = n_2 = 5$									
1.6	.487	.493	.584	.663	.780	.855	.914	.958	.976
	.558	.558	.641	.678	.772	.839	.897	.943	.963
1.4	.423	.442	.516	.610	.711	.809	.874	.918	.948
	.472	.484	.545	.610	.700	.789	.853	.900	.935
1.2	.321	.352	.432	.538	.635	.744	.821	.870	.907
	.354	.379	.440	.524	.615	.717	.788	.852	.893
1.0	.254	.271	.353	.424	.539	.646	.724	.787	.850
	.274	.281	.357	.411	.511	.617	.698	.762	.833
0.8	.195	.217	.267	.357	.452	.555	.635	.712	.774
	.197	.215	.254	.340	.432	.535	.610	.694	.773
0.6	.120	.149	.189	.263	.358	.433	.534	.598	.688
	.125	.147	.182	.259	.337	.420	.525	.609	.699
0.4	.088	.103	.142	.195	.259	.331	.428	.515	.589
	.083	.106	.136	.189	.254	.333	.439	.533	.633
0.2	.053	.068	.098	.139	.204	.277	.342	.445	.508
	.054	.069	.097	.140	.209	.287	.372	.484	.581
0.0	.052	.056	.088	.126	.190	.263	.332	.395	.486
	.054	.054	.090	.129	.192	.275	.365	.455	.550



$\mu_2$	$\mu_1$								
	0.0	0.2	0.4	0.6	0.8	1.0	1.2	1.4	1.6
$\alpha = .01, n_1 = n_2 = 5$									
1.6	.191	.205	.281	.339	.469	.578	.645	.757	.801
	.216	.217	.286	.336	.450	.441	.607	.717	.771
1.4	.143	.158	.226	.296	.391	.496	.572	.670	.727
	.151	.167	.227	.287	.370	.463	.535	.631	.683
1.2	.099	.122	.170	.247	.315	.404	.515	.578	.653
	.106	.128	.167	.228	.283	.381	.481	.541	.621
1.0	.070	.080	.121	.172	.231	.320	.414	.463	.555
	.070	.080	.115	.159	.210	.297	.397	.429	.527
0.8	.047	.058	.084	.135	.180	.256	.305	.391	.460
	.048	.059	.081	.124	.160	.230	.283	.373	.438
0.6	.032	.045	.054	.081	.132	.171	.228	.286	.353
	.034	.048	.052	.076	.120	.161	.210	.272	.348
0.4	.019	.024	.035	.050	.089	.127	.160	.203	.274
	.018	.023	.035	.048	.081	.120	.156	.206	.281
0.2	.009	.015	.027	.038	.058	.086	.124	.157	.203
	.009	.015	.024	.036	.055	.090	.122	.167	.220
0.0	.009	.011	.019	.032	.050	.073	.109	.143	.172
	.009	.012	.021	.032	.048	.081	.115	.152	.200

$\mu_1$									
$\mu_2$	0.0	0.2	0.3	0.4	0.5	0.7	0.9	1.0	
$\alpha = .05, n_1 = 10, n_2 = 10$									
1.0	.575 .651	.595 .761	.612 .676	.675 .720	.737 .755	.790 .811	.904 .898	.952 .948	.963 .958
0.9	.487 .550	.518 .579	.546 .599	.592 .622	.683 .705	.732 .735	.853 .836	.927 .917	.954 .950
0.7	.339 .374	.353 .383	.387 .410	.451 .466	.539 .535	.598 .598	.754 .737	.850 .845	.895 .894
0.5	.199 .211	.201 .220	.240 .249	.292 .294	.362 .358	.440 .429	.808 .601	.721 .731	.789 .807
0.4	.151 .154	.165 .169	.183 .180	.237 .233	.287 .282	.358 .347	.502 .509	.663 .676	.726 .751
0.3	.100 .106	.121 .122	.143 .148	.175 .170	.228 .226	.297 .300	.458 .468	.597 .630	.666 .709
0.2	.080 .077	.075 .075	.102 .102	.143 .142	.181 .183	.240 .244	.388 .413	.543 .597	.626 .688
0.1	.056 .058	.062 .062	.078 .078	.108 .105	.146 .151	.211 .224	.360 .393	.534 .596	.573 .642
0.0	.052 .049	.052 .054	.071 .066	.101 .105	.148 .156	.203 .219	.338 .378	.495 .558	.557 .631

\*Lower elements correspond to power of Fisher's method  
Upper elements correspond to power of Logit method.



REFERENCES

- Bahadur, R.R. (1971) Some limit theorems in Statistics, Society for Industrial and Applied Mathematics, Philadelphia PA.
- Berkson, J. (1944) Applications of the Logistic function to bioassay, Journal of the American statistical Association, 39, 357-365.
- Birnbaum, A (1954). Combining independent tests of significance, Journal of the American Statistical Association, 49, 559-575.
- Birnbaum, A (1955) Characterizations of complete classes of tests of some multiparameter hypotheses with applications to likelihood ratio tests, Annals of Mathematical Statistics, 26, 21-36.
- Brown, L.D., Cohen, A., and Strawderman, W.E. (1976). A complete class theorem for strict monotone likelihood ratio with applications, Annals of Statistics, 4, 712-722.
- Fisher, R.A. (1932). Statistical Methods for research workers, Oliver Boyd, Edinburg, London 4th Edition.
- George, E.O. and Mudholkar, G.S. (1977) University of Rochester Technical Report.
- Liptak, T., (1958). On the combination of independent tests, Magyar Tud Akad. Mat. Kutato Int. Kozl 3, 171-197.

Littell, R.C., and Folks, J.L. (1971) Asymptotic optimality of Fisher's method of combining independent tests, Journal of the American Statistical Association, 66, 802-804.

Littell, R.C., and Folks, J.L. (1973) Asymptotic optimality of Fisher's method of combining independent tests II, Journal of the American Statistical Association, 68, 193-194.

Mudholkar, G.S. (1969). A generalized monotone character of d.f.'s and moments of statistics from some well-known populations, Annals of the Institute of Statistical Mathematics, 21, 277-285.

Oosterhoff, J. (1969). Combination of one-sided statistical tests, Mathematisch Centrum Amsterdam.

Pearson, E.S. (1938). The probability integral transformation for testing goodness of fit and combining independent tests of significance, Biometrika, 30, 134-138.

Pearson, K. (1933). On a method of determining whether a sample of a given size  $n$  supposed to have been drawn from a parent population having known probability integral has been probably drawn at random, Biometrika, 25, 379-410.

Tippett, L.H.C. (1931). The method of statistics, Williams and Norgate, London, First Edition.

van Zwet, W.R., and Oosterhoff, J. (1967). On the combination of independent statistics, Annals of Mathematical Statistics, 38, 659-680.

Wallis, W.A. (1942). Compounding probability from independent significance tests, *Econometrica*, 10, 229-248.

Wiseand, H.S. (1976). A condition under which the Pitman and Bahadur approaches to efficiency coincide, *Annals of statistics*, 4, 1003-1011.

Wilkinson, B. (1951). A statistical consideration in psychological research, *Psychology Bulletin* 48, 156-156.



18 REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER <b>AFOSR/TR-79-0742</b>	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) <b>THE LOGIT METHOD FOR COMBINING TESTS</b>	5. TYPE OF REPORT & PERIOD COVERED <b>9 Interim rept.</b>	
7. AUTHOR(s) <b>E. OLUSEGUN/GEORGE GOVIND S./MUDHOLKAR</b>	6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Statistics University of Rochester Rochester, NY 14626	8. CONTRACT OR GRANT NUMBER(s) <b>15</b> ✓ <b>AFOSR-77-3260</b>	
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research/NM Bolling AFB, Washington, D.C. 20332	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS <b>16</b> 61102F 2304/A5	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) <b>12</b> 26p.	12. REPORT DATE <b>11</b> 1979	
	13. NUMBER OF PAGES <b>17</b> A5 25	
	15. SECURITY CLASS. (of this report) UNCLASSIFIED	
15a. DECLASSIFICATION/DOWNGRADING SCHEDULE		
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of this abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Combining Tests, Logit, Fisher's Omnibus procedure, Bahadur Efficiency, Admissibility.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A new combination statistic, the sum of the logits of P-values is introduced and its standardized form is found to be essentially equivalent to a standardized Student t in null distribution. Comparisons with Fisher's omnibus procedure by Bahadur ARE, membership in a complete class of tests and a Monte Carlo simulation of the power functions, show that both procedures are equivalent by the first criterion and neither procedure is generally dominant by either of the two criteria.		

410 276 Lm